

On the chromatic numbers of small-dimensional Euclidean spaces

A.B. Kupavskii^{1,2}

*Department of Number Theory, Mechs and Maths Faculty
Moscow State University
Moscow, Russia*

A.M. Raigorodskii^{3,4}

*Department of Mathematical Statistics, Mechs and Maths Faculty
Moscow State University
Moscow, Russia*

Abstract

In this paper, we will give a short survey concerning estimates on the chromatic numbers of Euclidean spaces in small dimensions. We will also present some new important bounds.

Keywords: Chromatic numbers, distance graphs.

¹ This work is supported by the grant 09-01-00294 of the Russian Foundation for Basic Research.

² E-mail: kupavskii@yandex.ru.

³ This work is supported by the grant 09-01-00294 of the Russian Foundation for Basic Research, by the grant MD-5414.2008.1 of the Russian President, by the grant NSh-691.2006.1 of the Leading Scientific Schools of Russia, and by the grant of "Dynastia" foundation.

⁴ E-mail: mraigor@yandex.ru.

1 Introduction

This paper is concerned with the classical Nelson – Erdős – Hadwiger problem on painting the space \mathbb{R}^n using the smallest possible number $\chi(\mathbb{R}^n)$ of colours in such a way that no two monochromatic points are at the distance 1 apart. In other words,

$$\chi(\mathbb{R}^n) = \min \left\{ \chi : \mathbb{R}^n = V_1 \sqcup \dots \sqcup V_\chi, \forall i \forall \mathbf{x}, \mathbf{y} \in V_i \quad |\mathbf{x} - \mathbf{y}| \neq 1 \right\}.$$

The quantity $\chi(\mathbb{R}^n)$ is called *the chromatic number of the space*.

Almost obviously, $\chi(\mathbb{R}^1) = 2$. However, even in the planar case, the problem still remains open. We only know that $4 \leq \chi(\mathbb{R}^2) \leq 7$ (see [1], [2]).

During the last decades, various upper and lower bounds have been obtained for the chromatic numbers of different spaces. In the next section, we will try to present the most recent results of this kind. In the last section, we will give our new improvements on those estimates.

2 An overview

First of all, we note that the problem we discuss here admits a natural reformulation in terms of graph theory. Indeed, we may consider the graph $\mathcal{G} = (\mathcal{V}, \mathcal{E})$, where

$$\mathcal{V} = \mathbb{R}^n, \quad \mathcal{E} = \{\{\mathbf{x}, \mathbf{y}\} : |\mathbf{x} - \mathbf{y}| = 1\}.$$

So the usual chromatic number $\chi(\mathcal{G})$ coincides with the value $\chi(\mathbb{R}^n)$. This interpretation suggests the study of colourings of *distance graphs*, i.e., of arbitrary subgraphs G in \mathcal{G} . All the best known lower bounds on $\chi(\mathbb{R}^n)$ are due to appropriate constructions of distance graphs.

For the three-dimensional space, the estimate $\chi(\mathbb{R}^3) \geq 5$ had been remaining unimproved for many years until in 2002 O. Nechushtan succeeded in replacing it by the bound $\chi(\mathbb{R}^3) \geq 6$ (see [3]). The tightest upper estimate was given in 2000 and published in 2002 by D. Coulson (see [4]): $\chi(\mathbb{R}^3) \leq 15$.

In four dimensions, the bound $\chi(\mathbb{R}^4) \geq 7$ was independently proposed in 1996 by K. Cantwell (see [5]) and in 2006 by L. Ivanov (see [6]). The last construction is a bit simpler than the first one. The upper estimate $\chi(\mathbb{R}^4) \leq 49$ was announced by Coulson, but we could find in the literature only the bound $\chi(\mathbb{R}^4) \leq 54$ (see [7]).

Starting from $n = 5$, we do not know any non-trivial upper estimates for the chromatic number. Certainly, $\chi(\mathbb{R}^n) \leq (\sqrt{n} + 1)^n$, which can be done by dividing the space into translates of a multicoloured cube. Also, we may cite D. Larman and C.A. Rogers' paper [8], where $\chi(\mathbb{R}^n) \leq (3 + o(1))^n$ is proved. However, nothing more precise and concrete has been done. So we just exhibit below a list of lower bounds for $5 \leq n \leq 12$:

- $\chi(\mathbb{R}^5) \geq 9$ (Cantwell, 1996, see [5]);
- $\chi(\mathbb{R}^6) \geq 11$ (J. Cibulka, 2008, see [9]);
- $\chi(\mathbb{R}^7) \geq 15$ (A.M. Raigorodskii, 2001, see [10]);
- $\chi(\mathbb{R}^8) \geq 16$ (see [11]);
- $\chi(\mathbb{R}^9) \geq 16$ (see [11]);
- $\chi(\mathbb{R}^{10}) \geq 19$ (Larman and Rogers, 1972, see [8]);
- $\chi(\mathbb{R}^{11}) \geq 20$ (Raigorodskii, 2001, see [10]);
- $\chi(\mathbb{R}^{12}) \geq 24$ (Larman and Rogers, 1972, see [8]).

As for growing dimensions, we have

$$(1.239 + o(1))^n \leq \chi(\mathbb{R}^n) \leq (3 + o(1))^n.$$

The lower bound is given in [10] and the upper one in [8].

For further references see [10], [11], [12].

3 New Results

First of all, we managed to achieve considerable improvements on lower bounds in the dimensions 9 – 12.

Theorem 1. *The estimates hold $\chi(\mathbb{R}^9) \geq 21$, $\chi(\mathbb{R}^{10}) \geq 23$, $\chi(\mathbb{R}^{11}) \geq 23$, $\chi(\mathbb{R}^{12}) \geq 25$.*

The results are obtained using a new series of distance graphs in the above-mentioned spaces.

At the same time, we propose a new general approach on "lifting" a bound in a dimension to a greater bound in a higher dimension:

Theorem 2.

References

- [1] Moser, L., Moser, W., *Solution to problem 10*, Canad. Math. Bull., **4** (1961), 187 - 189.
- [2] Hadwiger, H., *Ungelöste Probleme N 40*, Elemente der Math., **16** (1961), 103 - 104.
- [3] Nechushtan O., *On the space chromatic number*, Discrete Math., **256** (2002), 499 - 507.
- [4] Coulson D., *A 15-colouring of 3-space omitting distance one*, Discrete Math., **256** (2002), 83 - 90.
- [5] Cantwell, K., *Finite Euclidean Ramsey theory*, J. Comb. Theory, Ser. A, **73** (1996), 273 - 285.
- [6] Ivanov, L., *An estimate for the chromatic number of the space \mathbb{R}^4* , Russian Math. Surveys, **61** (2006), 371 - 372.
- [7] Radoičić, R., Tóth, G., *Note on the chromatic number of the space*, Aronov, B. (ed.) et al., Discrete and computational geometry, The Goodman-Pollack Festschrift, Berlin, Springer, Algorithms Comb., **25** (2003), 696 - 698.
- [8] Larman D.G., Rogers C.A., *The realization of distances within sets in Euclidean space*, Mathematika, **19** (1972), 1 - 24.
- [9] Cibulka, J., *On the chromatic number of real and rational spaces*, Geombinatorics, **18** (2008), 53 - 66.
- [10] Raigorodskii, A. M., *The Borsuk problem and the chromatic numbers of some metric spaces*, Russian Math. Surveys, **56** (2001), 103 - 139.
- [11] Székely, L., *Erdős on unit distances and the Szemerédi – Trotter theorems*, Paul Erdős and his Mathematics, Bolyai Series Budapest, J. Bolyai Math. Soc., Springer, **11** (2002), 649 - 666.
- [12] Brass, P., Moser W., and J. Pach, “Research problems in discrete geometry,” Springer, Berlin, 2005.